



Non-Native Invasive Plant Species Control Treatments

Timing, Methods and Herbicide Rates

Forestry Topic 31

www.dof.virginia.gov

March 2017

This is a list of management tactics for major invasive plants, not a comprehensive control plan. For recommendations specific to your property, consult a professional forester or land resource manager. Follow all label prohibitions, precautions and safety requirements during herbicide transport, storage, mixing and application.

INVASIVE PLANT	CONTROL TIMING	CONTROL METHOD	HERBICIDE RATES	NOTES
GRASSES	Japanese Stiltgrass	Prior to seed maturity	Manual – hand pull	Remove all roots
		Prior to seed maturity	Mechanical – mow/cut repeatedly	To reduce seed formation
		Early summer	Foliar spray	Glyphosate 0.1%
		Late spring – late summer	Foliar spray	Sethoxydim 1.5% or Glyphosate 0.5%-2%
	Wavyleaf Grass	Prior to seed maturity	Manual – pull small areas	Follow-up treatment required
		Late spring – fall	Foliar spray	Glyphosate 2%
		April – June	Foliar spray	Sethoxydim or Clethodim 1%
HERBS/FORBS	Garlic Mustard	Late spring – early summer, prior to any seedpods maturing to brown	Manual – hand pull and remove taproot	Do not leave flowering plants on ground, seeds will form; bag/ remove flowering plants
		March – June, before seedpods	Mechanical – mow or cut	To reduce seed formation
		Late fall – winter is best, but susceptible any time	Foliar spray on evergreen leaves	Glyphosate or Triclopyr 2%
VINES	Gen. Recommendations for All Vines	All year when soil is moist	Manual – hand pull small vines	Remove as many roots as possible to prevent resprouts
		Any	Manual or mechanical – cut to “treatable” height	Follow-up with foliar herbicide applied to resprouts
		June – October, through winter for evergreen species	Foliar spray	Glyphosate 2%-3% or Triclopyr 2%-5%
		June – February	Basal spray	Triclopyr ester 20%-25%
		June – August	Foliar spray	Metsulfuron 2-4 oz./acre
		June – February, late summer – fall ideal	Cut stump	Glyphosate or Triclopyr ester 20%-25%
	Japanese Honeysuckle	Before seed formation	Foliar spray	Glyphosate 2%-3% or Triclopyr 2%-5%

Non-Native Invasive Plant Species Control Treatments

Forestry Topic 31

www.dof.virginia.gov

March 2017


INVASIVE PLANT		CONTROL TIMING	CONTROL METHOD	HERBICIDE RATES	NOTES
VINES, continued	Porcelain-berry	June to October, late summer – early fall ideal	Foliar spray	Triclopyr 2%-3%	Manual ineffective due to extensive root system
	Oriental Bittersweet	All year, late summer – fall ideal	Injection or hack-n-squirt	Triclopyr amine or Glyphosate undiluted	Vines more than 1 inch in diameter
	Mile-a-Minute	May – October	Manual/mechanical – hand pull, mow or cut repeatedly		Protect skin from thorns
		May – July	Foliar spray	Glyphosate 1% or Triclopyr 1%-2%	Likely to injure other plants
	Kudzu	All year	Manually remove all root crowns		
		All year	Mechanical – mow and cover with plastic sheeting		Leave sheeting in place two years
		July – September	Mechanical – cut or mow to ground		Many, many years needed
		June – October	Foliar spray	Picloram 3%	*Restricted use pesticide
		June – October	Foliar spray	Metsulfuron 3-4 oz./acre, Triclopyr 4%, Clopyralid 1.3 pt./acre, Aminopyralid 7 oz./acre	Repeat in successive years
		June – February	Basal spray	Triclopyr ester 20%	Woody stems
		June – February	Injection or hack-n-squirt	Imazapyr, Triclopyr amine or Glyphosate undiluted	Vines more than 1 inch in diameter
SHRUBS	Gen. Recommendations for All Shrubs	When soil is moist	Manual – hand pull small plants		Roots left in ground resprout
		When fruit is not present	Mechanical – cut or mow		Follow-up treatment required
		June – February	Foliar spray	Imazapyr or Dicamba 1% or Triclopyr 2%	Several years (as needed)
		June – February	Cut stump	Imazapyr 5%-10% or Glyphosate 20%	Selective, minimal herbicide
		June – February	Basal spray	Triclopyr ester 20%	
	Multiflora Rose	See General Recommendations for all shrubs			
		June – October	Foliar spray	Glyphosate 2%-4% or Triclopyr 1%	
	Autumn Olive	See General Recommendations for All Shrubs			
	Chinese Privet	See General Recommendations for All Shrubs			
TREES	Gen. Recommendations for All Trees	June-February	Injection or hack-n-squirt	Triclopyr or Imazapyr undiluted	Small to large trees
		July – February	Basal spray	Triclopyr ester 20%-25%	Saplings
	Tree-of-Heaven	See General Recommendations for All Trees			Follow-up usually required
		Summer – fall	Foliar spray	Triclopyr 2%	Seedlings, saplings, resprouts

Non-Native Invasive Plant Species Control Treatments

Forestry Topic 31

www.dof.virginia.gov

March 2017



This institution is an equal opportunity provider.